arXiv:1505.00060v1 [math.CO] 1 May 2015

On well-covered, vertex decomposable and Cohen-Macaulay graphs.

Iván D. Castrillón[*], Roberto Cruz and Enrique Reyes[**]

4 de mayo de 2015

Resumen

Let $G = (V, E)$ be a graph. If G is a König graph or G is a graph without 3-cycles and 5-cycle, we prove that the following conditions are equivalent: Δ_G is pure shellable, R/I_Δ is Cohen-Macaulay, G is unmixed vertex decomposable graph and G is well-covered with a perfect matching of König type $e_1, ..., e_g$ without square with two e_i's. We characterize well-covered graphs without 3-cycles, 5-cycles and 7-cycles. Also, we study when graphs without 3-cycles and 5-cycles are vertex decomposable or shellable. Furthermore, we give some properties and relations between critical, extendables and shedding vertices. Finally, we characterize unicyclic graphs with each one of the following properties: unmixed, vertex decomposable, shellable and Cohen-Macaulay.

Keywords: Cohen-Macaulay, well-covered, unmixed, vertex decomposable, shellable, König, girth, unicyclic.

1. Introduction

Let G be a simple graph (without loops and multiply edges) whose vertex set is $V(G) = \{x_1, ..., x_n\}$ and edge set $E(G)$. Let $R = k[x_1, ..., x_n]$ be a polynomial ring over a field k, the *edge ideal* of G, denoted by $I(G)$, is the ideal of R generated by all monomials $x_i x_j$ such that $\{x_i, x_j\} \in E(G)$. G is a *Cohen-Macaulay graph* if $R/I(G)$ is a Cohen-Macaulay ring (See [2], [20]). A subset F of $V(G)$ is a *stable set* or *independent set* if $e \nsubseteq F$ for each $e \in E(G)$. The cardinality of the maximum stable set is denoted by $\beta(G)$. G is called *well-covered* if every maximal stable set has the same cardinality. On the other hand, a subset D of $V(G)$ is a vertex cover of G if $D \cap e \neq \emptyset$ for every $e \in E(G)$. The number of vertices in a minimum vertex cover of G is called the *covering number* of G and it is denoted by $\tau(G)$. This number coincide with $\mathrm{ht}(I(G))$, the *height* of $I(G)$. If the minimal vertex covers have the same cardinality then G is called *unmixed* graph. Notice that, D

---

[*] Partially supported by CONACYT and ABACUS-CINVESTAV

[**] Partially supported by SNI and ABACUS-CINVESTAV

is a vertex cover if and only if $V(G) \setminus D$ is an stable set. Hence, $\tau(G) = n - \beta(G)$ and G is well-covered if and only if G is unmixed. The *Stanley-Reisner complex* of $I(G)$, denoted by Δ_G, is the simplicial complex whose faces are the stable sets of G. Recall that a simplicial complex Δ is called *pure* if every facets has the same number of elements. Thus, Δ_G is pure if and only if G is well-covered. The well-covered property has been studied for some families of graphs: graphs with girth at least 5 (in [6]), graphs without 4 or 5-cycles (in [7]), simplicial, chordal and circular graphs (in [15]), block-cactus graphs (in [16]). In this paper (Theorem 4.6) we characterize the well-covered graph without 3-cycles, 5-cycles and 7-cycles.

In section 2 we give some properties and relation between critical, shedding and extendable vertices. We use these result for to study the following properties: well-covered and vertex decomposable of G; shellability of Δ_G and Cohen-Macaulay property of $R/I(G)$. These properties have been studied in ([2], [3], [5], [10], [11], [13], [17], [19], [20], [22]). In general, we have that (see [17], [20])

vertex-decomposable $\Rightarrow$ shellable $\Rightarrow$ sequentially Cohen-Macaulay

It is know that if each chordless cycle of G has length 3 or 5 then, G is vertex decomposable. (see [22]). Now, in section 3 we characterize vertex decomposable graphs without 3-cycles and 5-cycles. This result generalize the criterion given in [19] for shellable graphs. Furthermore, we characterize shellable graphs with girth at least 11.

On the other hand, we have following implications:

Unmixed vertex decomposable $\Rightarrow$ pure shellable $\Rightarrow$ Cohen-Macaulay $\Rightarrow$ well-covered

The equivalence between Cohen-Macaulay property and unmixed vertex decomposable property has been studied for some families of graphs: bipartite graphs (in [5] and [10]); very well covered graphs (in [4] and [13]); graphs with girth at least 5, block-cactus and graphs without 4-cycles and 5-cycles (in [11]). In section 5 we prove the equivalences of these properties for König graphs and graphs without 3-cycles and 5-cycles. Furthermore, we prove theses properties are equivalent to the following condition: G is unmixed graph with a perfect matching $e_1 = \{x_1, y_1\}, ..., e_g = \{x_g, y_g\}$ of König type without squares with two e_i's.

Finally, in section 6 we characterize the unicyclic graph with each one of the following properties: vertex decomposable, shellable, Cohen-Macaulay and well-covered.

2. Critical, extendable and shedding vertices.

Let X be a subset of $V(G)$, the *subgraph induced* by X in G, denoted by $G[X]$ is the graph with vertex set X and whose edge set is $\{\{x, y\} \in E(G) \mid x, y \in X\}$. Furthermore, $G \setminus X$ denote the induced subgraph $G[V(G) \setminus X]$. Now, if $v \in V(G)$ then the set of *neighbours* of v is denoted by $N_G(v)$ and its closed neighbourhood is $N_G[v] = N_G(v) \cup \{v\}$. The *degree* of v in G is $\deg_G(v) = |N_G(v)|$.

The contraction of the some vertex x in G denoted by G/x, is equivalent to the induced subgraph $G \setminus N_G[x]$. If G is unmixed, shellable, sequentially Cohen-Macaulay or vertex decomposable, then $G \setminus N_G[x]$ is unmixed, shellable, sequentially Cohen-Macaulay or vertex decomposable, respectively. (See [1], [19])

Definition 2.1 *G is vertex decomposable if G is a totally disconnected graph or there is a vertex v such that*

(a) *$G \setminus v$ and $G \setminus N_G[v]$ are both vertex decomposable, and*

(b) *each stable set in $G \setminus N_G[v]$ is not a maximal stable set in $G \setminus v$.*

A *shedding vertex* of G is any vertex which satisfies the condition (b). Equivalently, a shedding vertex is a vertex that satisfies: For every stable set S contained in $G \setminus N_G[v]$, there is some $x \in N_G(v)$ such that $S \cup \{x\}$ is stable.

Lemma 2.2 *If x is a vertex of G, then x is a shedding vertex if and only if $|N_G(x) \setminus N_G(S)| \geq 1$ for every stable set S of $G \setminus N_G[x]$.*

Proof. $\Rightarrow$) We take a stable set S of $G \setminus N_G[x]$. Since x is a shedding vertex then there is a vertex $z \in N_G(x)$ such that $S \cup \{z\}$ is stable set of $G \setminus x$. Thus, $z \notin N_G[S]$. Therefore, $|N_G(x) \setminus N_G(S)| \geq 1$.

$\Leftarrow$) We take a stable set S of $G \setminus N_G[x]$. Thus, there exists a vertex $z \in N_G(x) \setminus N_G(S)$. Since $z \in N_G(x)$, we have that $z \notin S$. Furthermore, $z \notin N_G(S)$ then, $S \cup \{z\}$ is a stable set of $G \setminus x$. Consequently, S is not a maximal stable set of $G \setminus x$. Therefore, x is a shedding vertex. $\square$

Definition 2.3 *Let S be a stable set of G. If x is of degree zero in $G \setminus N_G[S]$, then x is called isolated vertex in $G \setminus N_G[S]$, or we say that S isolates to x.*

By Lemma 2.2 we have that x is not a shedding vertex if and only if there exists a stable set S of $G \setminus N_G[x]$ such that $N_G(x) \subseteq N_G(S)$, i.e. x is an isolated vertex in $G \setminus N_G[S]$.

Corollary 2.4 *Let S be a stable set of G. If S isolates to x in G then x is not a shedding vertex in $G \setminus N_G[y]$ for all $y \in S$.*

Proof. Since S isolates to x then, S is a stable set in G and $\deg_{G\setminus N_G[S]}(x)=0$. Thus, $N_G(x)\subseteq N_G[S]\setminus S$Hence, if $y\in S$ and $G'=G\setminus N_G[y]$ then $x\in V(G')$ and $S\cap N_G[x]=\emptyset$. Consequently, $S'=S\setminus y$ is a stable set in $G'\setminus N_{G'}[x]$. Now, if $a\in N_{G'}(x)$ then there exists $s\in S'$ such that $\{a,s\}\in E(G')$. This implies $|N_{G'}(x)\setminus N_{G'}(S')|=0$. Therefore, by Lemma 2.2, x is not a shedding vertex in G'. $\square$

Theorem 2.5 *If x is a shedding vertex of G then one of the following condition holds:*

(a) *There is $y\in N_G(x)$ such that $N_G[y]\subseteq N_G[x]$.*

(b) *x is in a 5-cycle.*

Proof. We take $N_G(x)=\{y_1,y_2,...,y_k\}$. If there is a vertex y_i such that $N_G[y_i]\subseteq N_G[x]$ then G satisfies (a). On the other hand, there are $\{z_1,...,z_k\}\subseteq V(G)\setminus N_G[x]$ such that $\{y_i,z_i\}\in E(G)$ for $i\in\{1,...,k\}$. We can suppose that $L=\{z_1,z_2,...,z_q\}=\{z_1,...,z_k\}$ with $z_i\neq z_j$ for $1\leq i<j\leq q$. By Lemma 2.2 if L is a stable set of G then $|N_G(x)\setminus N_G(L)|\geq 1$. But $N_G(x)=\{y_1,...,y_k\}\subseteq N_G(L)$ then L is not a stable set. Hence, $k\geq 2$ and there exist $z_{i_1},z_{i_2}\in L$ such that $\{z_{i_1},z_{i_2}\}\in E(G)$. Thus, there exist y_{j_1} and y_{j_2} such that $y_{j_1}\neq y_{j_2}$ and $\{y_{j_1},z_{i_1}\},\{y_{j_2},z_{i_2}\}\in E(G)$. Therefore, $(x,y_{j_1},z_{i_1},z_{i_2},y_{j_2})$ is a 5-cycle of G. $\square$

Definition 2.6 *A vertex v is called simplicial if the induced subgraph $G[N_G(v)]$ is a complete graph or clique. Equivalently, a simplicial vertex is a vertex that appears in exactly one clique.*

Remark 2.7 *If $v,w\in V(G)$ such that $N_G[v]\subseteq N_G[w]$ then w is a shedding vertex of G* (see Lemma 6 in [22])*. In particular, if v is a simplicial vertex then any $w\in N_G(v)$ is a shedding vertex* (see Corollary 7 in [22])*.*

Corollary 2.8 *Let G be graph without 4-cycles. If x is a shedding vertex x of G then x is in a 5-cycle or there exists a simplicial vertex z such that $\{x,z\}\in E(G)$ with $|N_G[z]|\leq 3$.*

Proof. By Theorem 2.5, if x is not in a 5-cycle then there is $z\in N_G(x)$ such that $N_G[z]\subseteq N_G[x]$. If $\deg_G(z)=1$ then z is a simplicial vertex. If $\deg_G(z)=2$ then $N_G(z)=\{x,w\}$. Consequently, (z,x,w) is a 3-cycle and z is a simplicial vertex. Now, if $\deg_G(z)\geq 3$ then there are $w_1,w_2\in N_G(z)\setminus x$. Thus, $w_1,w_2\in N_G(x)$ since $N_G[z]\subseteq N_G[x]$. Hence, (w_1,z,w_2,x) is a 4-cycle of G. This is a contradiction. Therefore, $|N_G[z]|\leq 3$ and z is a simplicial vertex. $\square$

Remark 2.9 *If G is a 5-cycle with $V(G)=\{x_1,x_2,x_3,x_4,x_5\}$ then x_i is a shedding vertex. We can assume that $i=1$, then $\{x_3\}$ and $\{x_4\}$ are the stable sets in $G\setminus N_G[x_1]$. Furthermore, $\{x_3,x_5\}$ and $\{x_2,x_4\}$ are stable sets in $G\setminus x_1$. Hence,* *each stable set of* $G \setminus N_G[x_1]$ *is not a maximal stable set in* $G \setminus x_1$*. Therefore,* x_1 *is a shedding vertex.*

Definition 2.10 *A vertex* v *of* G *is critical if* $\tau(G \setminus v) < \tau(G)$*. Furthermore,* G *is called a vertex critical graph if each vertex of* G *is critical.*

Remark 2.11 *If* $\tau(G \setminus v) < \tau(G)$ *then* $\tau(G) = \tau(G \setminus v) + 1$*. Moreover,* v *is a critical vertex if and only if* $\beta(G) = \beta(G \setminus v)$*.*

Proof. If t is a minimal cover vertex such that $|t| = \tau(G \setminus v)$ then $t \cup v$ is a vertex cover of G. Thus, $\tau(G) \leq |t \cup v| = \tau(G \setminus v) + 1$. But, if $\tau(G) \geq \tau(G \setminus v) + 1$ then $\tau(G) = \tau(G \setminus v) + 1$.

Now, we have that $\tau(G)+\beta(G) = |V(G)| = |V(G \setminus v)|+1 = \tau(G \setminus v)+\beta(G \setminus v)+1$. Hence, $\beta(G) = \beta(G \setminus v)$ if and only if $\tau(G) = \tau(G \setminus v)+1$. Therefore, v is a critical vertex if and only if $\beta(G) = \beta(G \setminus v)$. ☐

Definition 2.12 *A vertex* v *of* G *is called an extendable vertex if* G *and* $G \setminus v$ *are well-covered graphs such that* $\beta(G) = \beta(G \setminus v)$*.*

Note that if v is an extendable vertex then every maximal stable set S of $G \setminus v$ contains a vertex of $N_G(v)$.

Corollary 2.13 *Let* G *be an unmixed graph and* $x \in V(G)$*. The following condition are equivalents:*

(a) *x is an extendable vertex.*

(b) $|N_G(x) \setminus N_G(S)| \geq 1$ *for every stable set* S *of* $G \setminus N_G[x]$*.*

(c) *x is a shedding vertex.*

(d) *x is a critical vertex and* $G \setminus x$ *is unmixed.*

Proof. $(a) \Leftrightarrow (b)$ ([6], Lemma 2).

$(b) \Leftrightarrow (c)$ By Lemma 2.2.

$(a) \Leftrightarrow (d)$ Since G is unmixed then by Remark 2.11, x is extendable if and only if x is a critical vertex and $G \setminus x$ is unmixed. ☐

3. Vertex decomposable and shellable properties in graphs without 3-cycles and 5-cycles.

Definition 3.1 *A 5-cycle* C *of* G *is called basic if* C *does not contain two adjacent vertices of degree three or more in* G*.*

Lemma 3.2 *If G is a graph then any vertex of degree at least 3 in a basic 5-cycle is a shedding vertex.*

Proof. Let $C = (x_1, x_2, x_3, x_4, x_5)$ be a basic 5-cycle. We suppose that $\deg_G(x_1) \geq 3$, since C is a basic 5-cycle then $\deg_G(x_2) = \deg_G(x_5) = 2$. We take a stable set S of $G \setminus N_G[x_1]$. Since $\{x_3, x_4\} \in E(G)$ then $|S \cap \{x_3, x_4\}| \leq 1$. If $x_3 \notin S$ or $x_4 \notin S$ then $S \cup \{x_2\}$ or $S \cup \{x_5\}$, respectively, is a stable set of $G \setminus x_1$. Therefore, x_1 is a shedding vertex. $\square$

Definition 3.3 *A simplicial complex Δ is shellable if the facets (maximal faces) of Δ can be ordered $F_1, ..., F_s$ such that for all $1 \leq i < j \leq s$, there exists some $v \in F_j \setminus F_i$ and some $l \in \{1, ..., j-1\}$ with $F_j \setminus F_l = \{v\}$. Furthermore, $F_1, ..., F_s$ is called a shelling of Δ. The facet set of Δ is denoted by $\mathcal{F}(\Delta)$. A graph G is called shellable if Δ_G is shellable.*

Proposition 3.4 *If G is shellable and x is a vertex of G then $G \setminus N_G[x]$ is shellable.*

Proof. ([19], Lemma 2.5). $\square$

Lemma 3.5 *Let G be a graph such that $\{z_1, ..., z_r\}$ is a stable set. If $N = \bigcup_{i=1}^{r} N_G[z_i]$ and $G^{i+1} = G^i \setminus N_{G^i}[z_i]$, with $G^1 = G$ then:*

(a) $\{z_{i+1}, ..., z_r\} \subseteq V(G^{i+1})$ *and*

(b) $G^{r+1} = G \setminus N$.

Proof. (a) By induction on i. If $i = 1$, since $\{z_1, ..., z_r\}$ is a stable set then $\{z_2, ..., z_r\} \cap N_G[z_1] = \emptyset$ and $\{z_2, ..., z_r\} \subseteq V(G \setminus N_G[z_1]) = V(G^2)$. Now, by induction hypothesis we have that $\{z_i, z_{i+1}, ..., z_r\} \subseteq V(G^i)$. Since $\{z_i, ..., z_r\}$ is a stable set then $\{z_{i+1}, ..., z_r\} \cap N_{G^i}[z_i] = \emptyset$. Hence, $\{z_{i+1}, ..., z_r\} \subseteq V(G^i \setminus N_{G^i}[z_i]) = V(G^{i+1})$.

(b) By induction on r. By $r = 1$, $N = N_G[z_1]$ then $G^2 = G \setminus N_G[z_1] = G \setminus N$. Now, if $r \geq 2$, we take $N^1 = \bigcup_{i=1}^{r-1} N_G[z_i]$ and by induction hypothesis we have that $G^r = G \setminus N^1$. By incise (a), we have that $\{z_r\} \subseteq V(G^r)$. We will prove that $N^1 \cup N_{G^r}[z_r] = N$. We have that $N^1 \subseteq N$. Furthermore, if $y \in N_{G^r}[z_r]$ then $\{y, z_r\} \in E(G^r)$. Thus, $\{y, z_r\} \in E(G)$ and $y \in N_G[z_r]$. Consequently, $y \in N$. Now, if $y \in N \setminus N^1$ then $y \in N_G[z_r]$ and $y \notin N_G[z_i]$ for $i \in \{1, ..., r-1\}$. This implies that $y, z_r \in V(G^r)$ and $\{y, z_r\} \in E(G^r)$ implying $y \in N_{G^r}[z_r]$. Hence, $N = N^1 \cup N_{G^r}[z_r]$. Therefore, $G^{r+1} = G^r \setminus N_{G^r}[z_r] = (G \setminus N^1) \setminus N_{G^r}[z_r] = G \setminus (N^1 \cup N_{G^r}[z_r]) = G \setminus N$. $\square$

Definition 3.6 *A subgraph H of G is called a c-minor (of G) if there exists a stable set S of G, such that $H = G \setminus N_G[S]$.*

Proposition 3.7 *Let H be a c-minor of G. If G is shellable then H is shellable.*

Remark 3.8 *In* ([21], Lemma 6) *is proved that: if G has a shedding vertex v and if both $G \setminus v$ and $G \setminus N_G[v]$ are shellables with shelling $F_1, ..., F_k$ and $G_1, ..., G_q$, respectively, then G is shellable. The order of the shelling is $F_1, ..., F_k, G_1 \cup \{v\}, ..., G_q \cup \{v\}$.*

Theorem 3.9 *Let G be a connected graph with a basic 5-cycle C. G is a shellable graph if and only if there is a shedding vertex $x \in V(C)$ such that $G \setminus x$ and $G \setminus N_G[x]$ are shellable graphs.*

Proof. $\Rightarrow$) We can suppose that $C = (x_1, x_2, x_3, x_4, x_5)$. If $G = C$ then G is shellable. By Remark 2.9 each vertex is a shedding vertex. Furthermore, $G_1 = G \setminus x_1$ is a path with shelling $\{x_2, x_4\}, \{x_2, x_5\}, \{x_3, x_5\}$ and $G \setminus N_G[x_1]$ is an edge. Therefore, G_1 and $G \setminus N_G[x_1]$ are shellable graphs. Now, we suppose $G \neq C$. We can assume that $\deg_G(x_1) \geq 3$ then, since C is a basic 5-cycle $\deg_G(x_2) = \deg_G(x_5) = 2$. Also, we can suppose $\deg_G(x_3) = 2$ and $\deg_G(x_4) \geq 2$. By Lemma 3.2, x_1 is a shedding vertex. $G \setminus N_G[x_1]$ is a shellable graph by Proposition 3.4. We will prove that G_1 is shellable. Since G is shellable then $G_2 = G \setminus N_G[x_2]$ is shellable. We assume that $F_1, ..., F_r$ is a shelling of Δ_{G_2}. Now, we take $G_3 = G \setminus N_G[x_3, x_5]$ then G_3 is shellable and whose shelling is $H_1, H_2, ..., H_k$. We take $F \in \mathcal{F}(\Delta_{G_1})$. If $x_2 \in F$ then $F \setminus x_2 \in \mathcal{F}(\Delta_{G_2})$ and there exists F_i such that $F = F_i \cup x_2$. If $x_2 \notin F$ then $x_3 \in F$ and $x_4 \notin F$. Thus, $x_5 \in F$. Hence, $F \setminus \{x_3, x_5\} \in \mathcal{F}(\Delta_{G_3})$ then there exist H_j such that $F = H_j \cup \{x_3, x_5\}$. This implies, $\mathcal{F}(\Delta_{G_1}) = \{F_1 \cup \{x_2\}, ..., F_r \cup \{x_2\}, H_1 \cup \{x_3, x_5\}, ..., H_k \cup \{x_3, x_5\}\}$. Furthermore, $F_1 \cup \{x_2\}, ..., F_r \cup \{x_2\}$ and $H_1 \cup \{x_3, x_5\}, ..., H_k \cup \{x_3, x_5\}$ are shellings. Now, $x_3 \in (H_j \cup \{x_3, x_5\}) \setminus (F_i \cup \{x_2\})$ and H_j is a stable set of G without vertices of C then $H_j \cup \{x_2, x_5\}$ is a maximal stable set of G since $N_G(x_2, x_5) = V(C)$ and $\{x_2, x_5\} \notin E(G)$. Consequently, $H_j \cup \{x_2, x_5\} = F_l \cup \{x_2\}$ for some $l \in \{1, ..., r\}$ then $(H_j \cup \{x_3, x_5\}) \setminus (F_l \cup \{x_2\}) = \{x_3\}$. Therefore, G_1 is a shellable graph.

$\Leftarrow$) By Remark 3.8. □

Definition 3.10 *If $v, w \in V(G)$ then the distance $d(u, v)$ between u and v in G is the length of the shortest path joining them, otherwise $d(u, v) = \infty$. Now, if $H \subseteq G$ then the distance from a vertex v to H is $d(v, H) = min\{d(v, u) \mid u \in V(H)\}$. Furthermore, we define $D_i(H) = \{v \in V(G) \mid d(v, H) = i\}$.*

Lemma 3.11 *Let G be a graph. If $W \subseteq V(G)$ then $D_0(W) = W$ and $D_i(W) = \{x \in V(G) \mid x \notin D_j(W)$ for $0 \leq j < i$ and $x \in N_G(D_{i-1}(W))\}$ for $i > 0$.*

Proof. We take $D_i = D_i(W)$. By definition $D_0 = \{v \in V(G) \mid d(v, W) = 0\} = W$. Now, we take $A = \{x \in V(G) \mid x \notin D_j \text{ for } 0 \leq j < i \text{ and } x \in N_G(D_{i-1})\}$. If $v \in D_i$ then $\mathrm{d}(v, W) = i$. Consequently, there is a vertex z such that $\{v, z\} \in E(G)$ and $\mathrm{d}(z, W) = i - 1$. Hence, $v \in N_G(D_{i-1}) \setminus N_G(D_j)$ for $0 \leq j < i$. Therefore, $v \in A$. On the other hand, $v \in A$ then $\mathrm{d}(v, W) \geq i$ since $v \notin D_j$ for

$0 \leq j < i$ and $\mathrm{d}(v, W) \leq i$ since $v \in N_G(D_{i-1})$. Therefore, $\mathrm{d}(v, W) = i$. $\square$

Definition 3.12 *A cut vertex of a graph is one whose removal increases the number of connected components. A block of a graph is a maximal subgraph without cut vertices. A connected graph without cut vertices with at least three vertices is called 2-connected graph.*

Remark 3.13 *Each connected component of a graph G is a c-minor of G.*

Definition 3.14 *A vertex of degree one is a leaf or free vertex. Furthermore, an edge which is incident with a leaf is called pendant.*

In the following result P is a property closed under c-minors, i.e., if G has the property P then each c-minor has the property P.

Theorem 3.15 *Let G be a graph without 3-cycles and 5-cycles with a 2-connected block B. If G satisfies the property P and B does not satisfy it then there exists $x \in D_1(B)$ such that* $\deg_G(x) = 1$

Proof. By contradiction, we assume that if $x \in D_1(B)$, then $|N_G(x)| > 1$. Thus, there exist $a, b \in N_G(x)$ with $a \neq b$. We can suppose that $a \in V(B)$. If $b \in V(B)$ then $G[\{x\} \cup V(B)]$ is 2-connected, but $B \subsetneq G[\{x\} \cup V(B)]$. This is a contradiction since B is a block. Consequently, $V(B) \cap N_G(x) = \{a\}$. Now, we suppose that $b \in D_1(B)$. Since there is not a 3-cycle in G, then $a \notin N_G(b)$. Hence, there exists $c \in N_G(b) \cap V(B)$ such that $c \neq a$. This implies $G[\{x, b\} \cup V(B)]$ is 2-connected. But $B \subsetneq G[\{x, b\} \cup V(B)]$, a contradiction then, $D_1(B) \cap N_G(x) = \emptyset$. Thus, $N_G(x) \cap (V(B) \cup D_1(B)) = \{a\}$ and $b \in D_2(B)$. Now, if $D_1(B) = \{x_1, \ldots, x_r\}$ then, there exist a_i such that $V(B) \cap N_G(x_i) = \{a_i\}$. Furthermore, there exists b_i such that $b_i \in N_G(x_i) \cap D_2(B)$. We can suppose that $L = \{b_1, \ldots, b_r\} = \{b_1, \ldots, b_s\}$ with $b_i \neq b_j$ for $1 \leq i < j \leq s$. We will prove that L is a stable set. Suppose that $\{b_i, b_j\} \in E(G)$, if $a_i = a_j$ then $(a_i, x_i, b_i, b_j, x_j, a_i)$ is a 5-cycle in G. This is a contradiction, consequently $a_i \neq a_j$ and the induced subgraph $G[\{x_i, b_i, b_j, x_j\} \cup V(B)]$ is 2-connected. But B is a block then, $\{b_i, b_j\} \notin E(G)$. Therefore, L is a stable set and $G' = G \setminus (N_G[L])$ satisfies the property P. Furthermore $D_1(B) \subset N_G(L)$ then, B is a connected component of G'. But, B does not satisfy P, this is a contradiction. Therefore, there exists a free vertex in $D_1(B)$. $\square$

Remark 3.16 *The following properties: unmixed, shellable, Cohen-Macaulay, sequentially Cohen-Macaulay and vertex decomposable are closed under c-minors.* (See [1], [20]).

Corollary 3.17 *Let G be a graph without 3-cycles and 5-cycles and B a 2-connected block. If G is shellable (unmixed, Cohen-Macaulay, sequentially Cohen-Macaulay or vertex decomposable) and B is not shellable (unmixed, Cohen - Macaulay, sequentially Cohen-Macaulay or vertex decomposable) then there exists*

$x \in D_1(B)$ such that $\deg_G(x) = 1$

Proof. From Remark 3.16 and Proposition 3.15 there exists $x \in D_1(B)$ such that $\deg_G(x) = 1$. ☐

Corollary 3.18 *Let G be a bipartite graph and B a 2-connected block. If G is shellable then there exists $x \in D_1(B)$ such that* $\deg_G(x) = 1$.

Proof. Since G is bipartite then B is bipartite. If H is a shellable bipartite graph then H has a free vertex. (see [19], Lemma 2.8). In particular, H is not 2-connected. Hence, B is not shellable. Therefore, by Corollary 3.17, there exists $x \in D_1(B)$ such that $\deg_G(x) = 1$. ☐

Lemma 3.19 *Let G be a graph without 3-cycles and 5-cycles. If G is vertex decomposable then G has a free vertex.*

Proof. Since G is vertex decomposable then, there is a shedding vertex x. Since there are not 5-cycles in G then by Theorem 2.5, there exists $y \in N_G[x]$ such that $N_G[y] \subseteq N_G[x]$. If $z \in N_G(y) \setminus x$ then (x, y, z) is a 3-cycle. This is a contradiction. Therefore, $N_G(y) = \{x\}$ and y is a free vertex. ☐

Remark 3.20 *If G is a vertex decomposable graph, then $G \setminus N_G[S]$ is a vertex decomposable graph for any S stable set of G*

Proof. See Theorem 2.5 in [1]. ☐

Theorem 3.21 *Let G be a graph without 3-cycles and 5-cycles and $\{x, y\} \in E(G)$ such that x is a free vertex. If $G_1 = G \setminus N_G[x]$ and $G_2 = G \setminus N_G[y]$, then G is vertex decomposable if and only if G_1 and G_2 are vertex decomposable.*

Proof. $\Rightarrow$) By Remark 3.20 G_1 and G_2 are vertex decomposable.

$\Leftarrow$) Since G_1 is vertex decomposable then $G_1 \cup \{x\} = G \setminus y$ is vertex decomposable. Furthermore, y is a shedding vertex by -Remark 2.7 and G_2 is vertex decomposable. Therefore, G is vertex decomposable. ☐

Corollary 3.22 *If G is a 2-connected graph without 3-cycles and 5-cycles then, G is not a vertex decomposable.*

Proof. If G is a vertex decomposable graph then, by Lemma 3.19, G has a free vertex. This is a contradiction since G is 2-connected. Therefore, G is not vertex decomposable. ☐

Definition 3.23 *Let G_1, G_2 be graphs. If $K = G_1 \cap G_2$ is a complete graph with $|V(K)| = k$ then $G = G_1 \cup G_2$ is called the k-clique-sum (or clique-sum) of G_1 and G_2 in K.*

Remark 3.24 *In* ([8], Proposition 4.1) *is proved that if C_n is a n-cycle then C_n is vertex decomposable, shellable or sequentially Cohen-Macaulay if and only* *if $n = 3$ or 5. Furthermore, a chordal graph is vertex decomposable.* (see [22], Corollary 7 (2))

Corollary 3.25 *If G is the 2-clique-sum of the cycles C_1 and C_2 with $|V(C_1)| = r_1 \leq r_2 = |V(C_2)|$ then, G is vertex decomposable if and only if $r_1 = 3$ or $r_1 = r_2 = 5$.*

Proof. $\Leftarrow$) First, we suppose that $r_1 = 3$. Consequently, we can assume $C_1 = (x_1, x_2, x_3)$ and $x_2, x_3 \in V(C_1) \cap V(C_2)$. Thus, x_1 is a simplicial vertex. Hence, by Remark 2.7 x_2 is a shedding vertex. Furthermore $G \setminus x_2$ and $G \setminus N_G[x_2]$ are trees then, $G \setminus x_2$ and $G \setminus N_G[x_2]$ are vertex decomposable graphs by Remark 3.24. Hence, G is vertex decomposable.

Now, we assume that $r_1 = r_2 = 5$ with $C_1 = (x_1, x_2, x_3, x_4, x_5)$ and $C_2 = (y_1, x_2, x_3, y_4, y_5)$. We take a stable set S in $G \setminus N_G[x_5]$. If $x_2 \in S$ then, $S \cup \{x_4\}$ is a stable set in $G \setminus x_5$. If $x_2 \notin S$ then, $S \cup \{x_1\}$ is a stable set in $G \setminus x_5$. Consequently, x_5 is a shedding vertex. Since x_2 is a neighbour of a free vertex in $G_1 = G \setminus x_5$ then, x_2 is a shedding vertex in G_1. Furthermore, since $G_1 \setminus x_2$ and $G_1 \setminus N_{G_1}[x_2]$ are forest then, they are vertex decomposable graphs by Remark 3.24. Thus, G_1 is vertex decomposable. Now, $G \setminus N_G[x_5] = C_2$ then it is vertex decomposable by Remark 3.24. Therefore, G is vertex decomposable.

$\Rightarrow$) By Corollary 3.22 we have that $\{r_1, r_5\} \cap \{3, 5\} \neq \emptyset$. If $r_1 \neq 3$ then $r_1 = 5$ or $r_2 = 5$. Consequently, we can assume that $\{C_1, C_2\} = \{C, C'\}$ where $C = (x_1, x_2, x_3, x_4, x_5)$ and $x_2, x_3 \in V(C) \cap V(C')$. Thus, $G \setminus N_G[x_5] = C'$ is vertex decomposable by Remark 3.20 then $|V(C')| \in \{3, 5\}$. But, $r_1 \neq 3$ then $|V(C')| = 5$ and $r_1 = r_5 = 5$. Therefore, $r_1 = 3$ or $r_1 = r_2 = 5$. □

Definition 3.26 *The girth of G is the length of the smallest cycle or if G is a forest we consider the girth of G infinite.*

Lemma 3.27 *Let G be a 2-connected graph with girth at least 11. Then G is not shellable.*

Proof. Let r be the girth of G, then there exists a cycle $C = (x_1, x_2, ..., x_r)$. If $G = C$ then, G is not shellable. Hence, $G \neq C$ then, $D_1(C) \neq \emptyset$. We take $y \in D_1(C)$, without loss of generality we can assume that $\{x_1, y\} \in E(G)$. If $\{x_i, y\} \in E(G)$ for some $i \in \{2, ..., r\}$ then we take the cycles $C_1 = (y, x_1, x_2, ..., x_i)$ and $C_2 = (y, x_1, x_r, x_{r-1}, ..., x_i)$. Thus, $|V(C_1)| = i + 1$ and $|V(C_2)| = r - i + 3$ Since r is the girth of G then $i + 1 \geq r$ and $r - i + 3 \geq r$. Consequently, $3 \geq i$ implies $4 \geq r$. But $r \geq 11$, this is a contradiction. This implies that $|N_G(y) \cap V(C)| = 1$. Now, we suppose that there exist $y_1, y_2 \in D_1(C)$ such that $\{y_1, y_2\} \in E(G)$. We can assume that $\{x_1, y_1\}, \{x_i, y_2\} \in E(G)$. Since $r \geq 11$ then, there are not 3-cycles in G. In particular, $x_1 \neq x_i$. Now, we take the cycle $C' = (y_1, x_1, ..., x_i, y_2)$ and $C'' = (y_1, x_1, x_r, x_{r-1}, ..., x_i, y_2)$. But, $|V(C')| = i + 2$

and $|V(C'')| = r - i + 4$. Since r is the girth then $i + 2 \geq r$ and $r - i + 4 \geq r$. Hence, $4 \geq i$ and $6 \geq r$. This is a contradiction then $D_1(C)$ is a stable set. Now, since G is 2-connected then for each $y \in D_1(C)$ there exists $z \in N_G(y) \cap D_2(C)$. Now, if there exist $z_1, z_i \in D_2(C)$ such that $\{z_1, z_i\} \in E(G)$ then, there exist $y_1, y_i \in D_1(C)$ such that $\{z_1, y_1\}, \{z_i, y_i\} \in E(G)$. Since there are not 3-cycles then $y_1 \neq y_i$. We can assume that $\{x_1, y_1\}, \{x_i, y_i\} \in E(G)$. Since there are not 5-cycles then $i \neq 1$. Consequently, there exist cycles $C'_1 = (x_1, ..., x_i, y_i, z_i, z_1, y_1)$ and $C'_2 = (x_i, ..., x_r, x_1, y_1, z_1, z_i, y_i)$. This implies $r \leq |V(C'_1)| = i + 4$ and $r \leq |V(C'_2)| = r - i + 6$. Hence, $i \leq 6$ implies $r \leq 10$. This is a contradiction then $D_2(C)$ is a stable set. Thus, C is a connected component of $G \setminus N_G[D_2(C)]$. But C is not shellable, therefore G is not shellable. $\square$

Theorem 3.28 *If G has girth at least 11 then G is shellable if and only if there exists $x \in V(G)$ with $N_G(x) = \{y\}$ such that $G \setminus N_G[x]$ and $G \setminus N_G[y]$ are shellables.*

Proof. $\Leftarrow$) By ([19], Theorem 2.9).

$\Rightarrow$) Since shellability is closed under c-minors, it is only necessary to prove that there exist $x \in V(G)$, such that $\deg_G(x) = 1$. If every block of G is an edge then G is a forest and there exist $x \in V(G)$ with $\deg_G(x) = 1$. Hence, we can assume that there exist B a 2-connected block of G. Since B is an induced subgraph of G, then its girth is at least 11. Thus, by Lemma 3.27, B is not shellable. Consequently, by Theorem 3.15, there exist $x \in D_1(B)$ such that $\deg_G(x) = 1$. $\square$

4. König graph and well-covered graphs without 3-cycles and 5-cycles.

In [4] and [13] proved that Cohen-Macaulay, pure shellable, unmixed vertex decomposable are equivalent properties if G is a very well-covered graph. Furthermore, they gave a theorem similar to theorem given in [10] by Herzog and Hibi for bipartite graphs.

In this paper we denoted by Z_G the set of the isolated vertices, that is,

$$Z_G = \{x \in V(G) \mid \deg_G(x) = 0\}.$$

Definition 4.1 *A graph G is called very well-covered if it is well-covered without isolated vertices and with $2ht(I(G)) = |V(G)|$.*

Definition 4.2 *G is a König graph if $\tau(G) = \nu(G)$ where $\nu(G)$ is the maximum number of pairwise disjoint edges. A perfect matching of König type of G is a collection $e_1, ..., e_g$ of pairwise disjoint edges whose union is $V(G)$ and $\tau(G) = g$.*

Proposition 4.3 *Let G be a König graph and $G' = G \setminus Z_G$. Then the following are equivalent:*

(a) *G is unmixed*

(b) *If $G' \neq \emptyset$, then G' has a perfect matching $e_1, ..., e_g$ of König type such that for any two edges $f_1 \neq f_2$ and for two distinct vertices $x \in f_1$, $y \in f_2$ contained in some e_i, one has that $(f_1 \setminus x) \cup (f_2 \setminus y)$ is an edge.*

Proof. By ([14], Lemma 2.3 and Proposition 2.9). □

Lemma 4.4 *G is an unmixed König graph if and only if G is totally disconnected or $G^{'} = G \setminus Z_G$ is very well-covered.*

Proof. $\Rightarrow$) If G is not totally disconnected, then from Proposition 4.3 G' has a perfect matching $e_1, ..., e_g$ of König type. Hence, $|V(G')| = 2\tau(G') = 2ht(I(G'))$. Furthermore, G' is unmixed then G' is well-covered.

$\Leftarrow$) If G is totally disconnected then $\nu(G) = 0$ and $\tau(G) = 0$. Hence, G is unmixed König graph. Now, if G is not totally disconnected then G' is very well-covered. Consequently, by ([9], Remark 2.2) G' has a perfect matching. Thus, $\nu(G') = |V(G')|/2 = ht(G') = \tau(G')$. Furthermore, G' is well-covered. Hence, G' is unmixed König graph. Therefore, G also is unmixed König graph. □

Lemma 4.5 *If G is a well-covered graph without 3-cycles, 5-cycles and 7-cycles, then G is a König graph.*

Proof. By induction on $|V(G)|$. We take $x \in V(G)$ then, by Remark 3.16 $G_1 = G \setminus N_G[x]$ is an unmixed graph. Furthermore, G_1 does not contain 3-cycles, 5-cycles and 7-cycles then, by induction hypothesis, G_1 is König, by Proposition 4.3. If $G_2 = G_1 \setminus Z_G = \emptyset$ then $V(G) = N_G[x] \cup Z_[G_1]$. Hence, $x \cup Z_{G_1}$ and $N_G(x)$ are stable set and G is bipartite. Consequently, G is König. Therefore, we can assume that $G_2 \neq \emptyset$ implying G_2 has a perfect matching $e_1 = \{x_1, y_1\}, ..., e_g = \{x_g, y_g\}$ of König type. We can assume that $N_G(x) = \{z_1, ..., z_r\}$ and $D = \{x_1, ..., x_g\}$ is a minimal vertex cover of G. Thus, $F = \{y_1, ..., y_g\}$ is a maximal stable set of G_2. Now, we take the subsets: $A_1 = N_G(z_1, ..., z_r) \cap D$, $B_1 = \{y_j \in F \mid x_j \in A_1\}$, $B_2 = N_G(z_1, ..., z_r) \cap F$ and $A_2 = \{x_j \in D \mid y_j \in B_2\}$. If there exists $y_i \in B_1 \cap B_2$ then $x_i \in A_1$ and there exist $z_k, z_p \in N_G(x)$ such that $\{x_i, z_k\}, \{y_i, z_p\} \in E(G)$. If $k = p$ then (z_k, x_i, y_i) is a 3-cycle and if $k \neq p$ then (x, z_k, x_i, y_i, z_p) is a 5-cycle. This is a contradiction, consequently $B_1 \cap B_2 = \emptyset$. Now, we take the subsets $B_3 = (N_G(A_2) \cap F) \setminus B_2$, $A_3 = \{x_j \in D \mid y_j \in B_3\}$, $B_4 = (N_G(A_1) \cap F) \setminus B_1$ and $A_4 = \{x_j \in D \mid y_j \in B_4\}$. If $y_i \in B_1 \cap B_3$ then there exist $x_j \in A_2$ and $z_k, z_p \in N_G(x)$ such that $\{x_i, z_k\}, \{y_i, x_j\}$ and $\{y_j, z_p\} \in E(G)$. Hence, if $k = p$ then $(z_k, x_i, y_i, x_j, y_j)$ is a 5-cycle and if $k \neq p$ then $(x, z_k, x_i, y_i, x_j, y_j, z_p)$ is a 7-cycle. A contradiction, implying $B_1 \cap B_3 = \emptyset$. Now, if $y_i \in B_2 \cap B_4$ then there exist $x_j \in A_1$ and $z_k, z_p \in N_G(x)$ such that $\{x_j, z_k\}, \{x_j, y_i\}$ and $\{y_i, z_p\} \in E(G)$. Consequently, if $k = p$ then (z_k, x_j, y_i) is a 3-cycle and if $k \neq p$ then (x, z_k, x_j, y_i, z_p) is a 5-cycle. This is a contradiction, hence $B_2 \cap B_4 = \emptyset$. Now, if $y_i \in B_3 \cap B_4$ then there exist $x_j \in A_1$, $x_q \in A_2$ and $z_k, z_p \in N_G(x)$ such that $\{x_j, y_i\}$, $\{x_q, y_i\}$, $\{x_j, z_k\}$ and $\{y_q, z_p\} \in E(G)$. Thus, if $k = p$ then $(z_k, x_j, y_i, x_q, y_q)$ is a 5-cycle and if $k \neq p$ then $(x, z_k, x_j, y_i, x_q, y_q, z_p)$ is a 7-cycle. This is a contradiction, implying $B_3 \cap B_4 = \emptyset$. Therefore, B_1, B_2, B_3, B_4, B_5 are pairwise disjoint sets, where $B_5 = F \setminus (B_1 \cup B_2 \cup B_3 \cup B_4)$.

Now, we will prove that if $A' = A_1 \cup A_4 \cup A_5$ and $B' = B_1 \cup B_4 \cup B_5$ then $N_G(B') \subseteq A'$. First, we observe that $N_G(Z_{G_1}) \subseteq N_G(x)$. Since $B_3 \cap B_5 = \emptyset$ then $N_G(B_5) \cap A_2 = \emptyset$. Furthermore, if $x_i \in N_G(B_5) \cap A_3$ then, there exist $y_j \in B_5$ and $x_q \in A_2$ such that $\{x_i, y_j\}$, $\{x_q, y_i\} \in E(G)$. Hence, since G_1 is unmixed and by Proposition 4.3 $\{x_q, y_j\} = (\{x_q, y_i\} \setminus y_i) \cup (\{x_i, y_j\} \setminus x_i) \in E(G)$. Consequently, $y_j \in B_3 \cap B_5$, this is a contradiction. Therefore, $N_G(B_5) \cap A_3 = \emptyset$ and $N_G(B_5) \subseteq A'$. Also, since $B_1 \cap B_3 = \emptyset$ then $N_G(B_1) \cap A_2 = \emptyset$. Furthermore, if $x_i \in N_G(B_1) \cap A_3$ then there exist $y_j \in B_1$, $x_q \in A_2$ such that $\{x_i, y_j\}, \{x_q, y_i\} \in E(G)$. Since G_1 is unmixed, by Proposition 4.3 $\{x_q, y_j\} = (\{x_i, y_j\} \setminus x_i) \cup (\{x_q, y_i\} \setminus y_i) \in E(G)$. Then $y_j \in B_1 \cap B_3$. This is a contradiction, thus $N_G(B_1) \cap A_3 = \emptyset$, and $N_G(B_1) \subseteq A'$. Now, since $B_3 \cap B_4 = \emptyset$ then $N_G(B_4) \cap A_2 = \emptyset$. Furthermore, if $x_i \in N_G(B_4) \cap A_3$ then there exist $y_j \in B_4$ and $x_q \in A_2$ such that $\{x_i, y_j\}$ and $\{x_q, y_i\} \in E(G)$. Consequently, since G_1 is unmixed $\{x_q, y_j\} = (\{x_i, y_j\} \setminus x_i) \cup (\{x_q, y_i\} \setminus y_i) \in E(G)$. This is a contradiction since $B_3 \cap B_4 = \emptyset$. Hence, $N_G(B_4) \cap A_3 = \emptyset$ and $N_G(B_4) \subseteq A'$. Therefore, $N_G(B') = A'$.

This implies $G_3 = G \setminus N_G[B'] = G \setminus (A' \cup B')$. Furthermore, since B' is a stable set then, G_3 is unmixed without 3-cycles, 5-cycles and 7-cycles. Now, if $G \neq G_3$ then, by induction hypothesis G_3 is König. That is, $\tau(G_3) = \nu(G_3)$. Consequently, if D' is a minimal vertex cover of G_3 then $D' \cup A'$ is a vertex cover of G since $N_G(B') = A'$. Furthermore, $G[A' \cup B']$ has a perfect matching with $|A'|$ elements. Thus, $\tau(G) \leq \tau(G_3) + |A'| = \nu(G_3) + |A'| \leq \nu(G)$. Hence, $\tau(G) = \nu(G)$, implying, G is König. Therefore, we can assume $G = G_3$.

Now, if there exist $x_i, x_j \in A_2$ such that $\{x_i, x_j\} \in E(G)$ then there exist $z_k, z_p \in N_G(x)$ such that $\{y_i, z_k\}$ and $\{y_j, z_p\} \in E(G)$. If $k = p$ then $(z_k, y_i, x_i, x_j, y_j)$ is a 5-cycle and if $k \neq p$ then $(x, z_k, y_i, x_i, x_j, y_j, z_p)$ is a 7-cycle. This is a contradiction, then A_2 is a stable set. Similarity, if $\{x_i, x_j\} \in E(G)$ with $x_i \in A_2$ and $x_j \in A_3$, then there exist $x_q \in A_2$ such that $\{x_q, y_j\} \in E(G)$. If $q = i$ then (x_i, x_j, y_j) is a 3-cycle. This is a contradiction, hence $q \neq i$. Consequently, since G_3 is unmixed and by Proposition 4.3, $\{x_i, x_q\} = (\{x_i, x_j\} \setminus x_j) \cup (\{x_q, y_j\} \setminus y_j) \in E(G_3)$. This is a contradiction since A_2 is a stable set of G. Thus, there are not edges between A_2 and A_3. Now, $\{x_i, x_j\} \in E(G)$ with $x_i, x_j \in A_3$ then there is a vertex $x_q \in A_2$ such that $\{x_q, y_i\} \in E(G)$. Hence, since G_3 is unmixed then $\{x_j, x_q\} = (\{x_i, x_j\} \setminus x_i) \cup (\{x_q, y_i\} \setminus y_i) \in E(G_3)$. But, there are not edges between A_2 and A_3. Therefore, A_3 is a stable set, implying $A_2 \cup A_3$ is also a stable set in G. This implies, $\{x\} \cup Z_{G_1} \cup B_2 \cup B_3$ and $N_G(x) \cup A_2 \cup A_3$ are stable sets, since $N_G(Z_{G_1}) \subseteq N_G(x)$, implying G is bipartite. Therefore, G is König. $\square$

Theorem 4.6 *If G is a graph without 3-cycles, 5-cycles and 7-cycles then the following conditions are equivalent:*

(1) *G is well-covered.*

(2) *G has a perfect matching $e_1, ..., e_g$ of König type such that if $f_1, f_2 \in E(G)$ with $a \in f_1$, $b \in f_2$ and $\{a, b\} = e_i$ then, $(f_1 \setminus a) \cup (f_2 \setminus b) \in E(G)$.*

Proof. (1) $\Rightarrow$ (2) By Lemma 4.5, G is König. Furthermore, since G is well-covered then by Proposition 4.3 G satisfies (2).

(2) $\Rightarrow$ (1) By Proposition 4.3, G is well covered. $\square$

Example 4.7 *The subgraph G is unmixed without 3-cycles and 5-cycles. If G is König then G has a perfect matching by* ([14], Lemma 2.3) *and $|V(G)|$ is even. But $|V(G)| = 9$, thus G is not König.*



Lemma 4.8 *If G is unmixed graph then for each $x \in V(G)$, $N_G(x)$ does not contain two free vertices.*

Proof. By contradiction suppose that there exists $x \in V(G)$ such that $y_1, ..., y_s$ are free vertices in $N_G(x_1)$ with $s \geq 2$. Hence, $G' = G \setminus N_G[y_1, ..., y_s] = G \setminus \{x, y_1, ..., y_s\}$ is unmixed. Now, we take S a maximal stable set of G'. Thus, $|S| = \beta(G')$ since G' is unmixed. Consequently, $S_1 = s \cup \{y_1, ..., y_s\}$ is stable set in G. Furthermore, we take S_2 a maximal stable in G such that $x \in S_2$. This implies that $S_2 \setminus x$ is a stable set in G'. Then, $|S_2| \leq \beta(G') + 1 \leq |S| + s = |S_1|$. This is a contradiction, since S_2 is a maximal stable set and G is unmixed. $\square$

Lemma 4.9 *Let G be an unmixed connected graph without 3-cycles and 5-cycles. If C is a 7-cycle and H is a c-minor of G with $C \subseteq H$ such that C has three non* *adjacent vertices of degree 2 in H then C is a c-minor of G.*

Proof. We take a minimal c-minor H of G such that $C \subseteq H$ and C has three non adjacent vertices of degree 2 in H. We can suppose that $C = (x, z_1, w_1, a, b, w_2, z_2)$ with $\deg_H(x) = \deg_H(w_1) = \deg_H(w_2) = 2$. If $\{z_1, b\} \in E(H)$ then, (z_1, b, w_2, z_2, x) is a 5-cycle of G. Thus, $\{z_1, b\} \notin E(H)$, similarity $\{z_2, a\} \notin E(H)$. Furthermore, since G does not have 3-cycles then $\{z_1, z_2\}, \{z_1, a\}$, $\{z_2, b\} \notin E(H)$. Hence, C is an induced cycle in H. On the other hand, if there exists $v \in V(H)$ such that $\mathrm{d}(v, C) \geq 2$, then $H' = H \setminus N_G[v]$ is a c-minor of G and $C \subseteq H'$. This is a contradiction by the minimality of H. Therefore, $\mathrm{d}(v, C) \leq 1$ for each $v \in V(H)$.

Now, if $\deg_H(b) \geq 3$ then there exists $c \in V(H) \setminus V(C)$ such that $\{b, c\} \in E(H)$. If $\{c, z_2\} \notin E(G)$ then $N_{H_1}(z_2)$ has two leaves w_2 and x in $H_1 = H \setminus N_H[w_1, c]$. This is a contradiction by Lemma 4.8. Thus, $\{c, z_2\} \in E(H)$. Furthermore, $\{a, c\}$, $\{z_1, c\} \notin E(H)$ since (a, b, c) and (z_1, w_1, a, b, c) are not cycles in G. Hence, if $\deg_H(c) \geq 3$ then there exists $d \in V(H) \setminus V(C)$ such that $\{c, d\} \in E(H)$. Also, $\{d, b\}, \{d, z_2\}, \{d, z_1\} \notin E(H)$ since (c, b, d), (z_2, b, c) and (z_1, x, z_2, c, d) are not cycles of G. But, $\mathrm{d}(d, C) \leq 1$ then, $\{a, d\} \in E(H)$. Consequently, $N_{H_2}(z_1)$ has two leaves w_1 and x in $H_2 = H \setminus N_H[d, w_2]$. This is a contradiction, by Lemma 4.8 then, $\deg_H(c) = 2$. But, $N_{H_3}(z_2)$ has two leaves w_2 and c in $H_3 = H \setminus N_H[a]$. A contradiction, therefore, $\deg_H(b) = 2$. Similarity, $\deg_H(a) = 2$.

Now, if $\deg_H(z_2) \geq 3$ then there exists $c' \in V(H) \setminus V(C)$ such that $\{c', z_2\} \in E(H)$. If there exists $d' \in V(H) \setminus V(C)$ such that $\{c', d'\} \in E(H)$ then, $\{d', z_1\}$ or $\{d', z_2\} \in E(G)$, since $\mathrm{d}(d', C) \leq 1$. But (c', d', z_2) and (x, z_2, c', d', z_1) are not cycles of H. Thus, $N_H(c') \subseteq \{z_1, z_2\}$. Consequently, $N_{H_4}(z_2)$ has two leaves x and c' in $H_4 = H \setminus N_H[w_1]$. This is a contradiction by Lemma 4.8, hence, $\deg_H(z_2) = 2$. Similarity, $\deg_H(z_1) = 2$. Furthermore, by the minimality, H is connected. Therefore, $H = C$ and C is a c-minor of G. □

5. König and Cohen-Macaulay graphs without 3-cycles and 5-cycles.

Proposition 5.1 *Let G be a König graph. If $G' = G \setminus Z_G$ then, the following properties are equiva-lent:*

(i) *G is unmixed vertex decomposable.*

(ii) *Δ_G is pure shellable.*

(iii) *$R/I(G)$ is Cohen-Macaulay.*

(iv) $G' = \emptyset$ *or* G' *is an unmixed graph with a perfect matching* $e_1, ..., e_g$ *of König type without squares with two* e_i*'s.*

(v) $G' = \emptyset$ *or there exists relabelling of vertices* $V(G') = \{x_1, ..., x_h, y_1, ..., y_h\}$ *such that* $\{x_1, y_1\}, ..., \{x_h, y_h\}$ *is a perfect matching,* $X = \{x_1, ..., x_h\}$ *is a minimal vertex cover of* G' *and the following conditions holds:*

   (a) *If* $a_i \in \{x_i, y_i\}$ *and* $\{a_i, x_j\}, \{y_j, x_k\} \in E(G')$*, then* $\{a_i, x_k\} \in E(G')$ *for distinct* i, j, k*;*

   (b) *If* $\{x_i, y_j\} \in E(G')$*, then* $i \leq j$*.*

Proof. (i)⇔(ii)⇔(iii) In each case G is unmixed and König then by Lemma 4.4, G is totally disconnected or G' is very well-covered. If G is totally disconnected then we obtain the equivalences. Now, if G' is very well covered then by ([13], Theorem 1.1) we obtain the equivalences.

(iv)⇔(iii) We can assume that $G' \neq \emptyset$ then by Lemma 4.4, G' is very well-covered. Hence, by ([4], Theorem 3.4) since $G = G' \cup Z_G$ therefore, G Cohen-Macaulay.

(iii)⇒ (v) Since $R/I(G)$ is Cohen-Macaulay then G is unmixed. Thus, by Lemma 4.4, we can assume that G' is very well-covered and by ([13], Lemma 3.1) G' satisfies (v).

(v)⇒(iv) We can assume that $G' \neq \emptyset$. Since, G' has a perfect matching $e_1, ..., e_h$ then $\nu(G') = h$. Furthermore, X is a minimal vertex cover, then $\tau(G') = h$ and $e_1, ..., e_h$ is a perfect matching of König type. Thus, from (a) and Proposition 4.3 G' is unmixed. Finally, from (b) there are not square with two e_i's. Since, $\{y_1, ..., y_h\}$ is a stable set. Therefore, G' satisfies (iv). □

Corollary 5.2 *Let* G *be a connected König graph. If* G *is a Cohen-Macaulay graph then* G *is an isolated vertex or* G *has at least a free vertex.*

Proof. By Proposition 5.1 (v), G is an isolated vertex or G has a perfect matching $e_1 = \{x_1, y_1\}, ..., e_h = \{x_h, y_h\}$ where $\{x_1, ..., x_h\}$ is a minimal vertex cover. Thus, $\{y_1, ..., y_g\}$ is a maximal stable set. Furthermore, if $\{x_i, y_j\}$ then $i \leq j$. Therefore, $N_G(y_1) = \{x_1\}$, that is, y_1 is a free vertex. □

Lemma 5.3 *Let* G *be an connected Cohen-Macaulay graph with a perfect matching* e_1, ..., e_g *of König type without square with two* e_i*'s and* $g \geq 2$*. For each* $z \in V(G)$ *we have that:*

(a) *If* $\deg_G(z) \geq 2$ *then there exist* $\{z, w_1\}, \{w_1, w_2\} \in E(G)$ *such that* $\deg_G(w_2) = 1$*. Furthermore,* $e_i = \{w_1, w_2\}$ *for some* $i \in \{1, ..., g\}$*.*

(b) *If* $\deg_G(z) = 1$ *then there exist* $\{z, w_1\}, \{w_1, w_2\}, \{w_2, w_3\} \in E(G)$ *such that* $\deg_G(w_3) = 1$*. Furthermore,* $e_i = \{z, w_1\}$ *and* $e_j = \{w_2, w_3\}$ *for some*

$i, j \in \{1, ..., g\}$.

Proof. Since $e_1 = \{x_1, y_1\}, ..., e_g = \{x_g, y_g\}$ is a perfect matching of König type we can assume $D = \{x_1, ..., x_g\}$ a minimal vertex cover. Thus, $F = \{y_1, ..., y_g\}$ is a maximal stable set. We take any vertex $z \in V(G)$. By Proposition 5.1 we can assume that if $\{x_i, y_j\} \in E(G)$ then $i \leq j$.

(a) First, we suppose that $z = x_k$ and there is a vertex x_j in $N_G(x_k)$. If y_j is a free vertex then, we take $w_1 = x_j$ and $w_2 = y_j$. Consequently, $e_j = \{w_1, w_2\}$. Now, we can assume $N_G(y_j) \setminus x_j = \{x_{p_1}, ..., x_{p_r}\}$ with $p_1 < \cdots < p_r < j$. If y_{p_1} is not a free vertex then, there is a vertex x_p with $p < p_1$ such that $\{x_p, y_{p_1}\} \in E(G)$. Since G is unmixed and from Proposition 4.3 we obtain that $\{x_p, y_j\} = (\{x_p, y_{p_1}\} \setminus y_{p_1}) \cup (\{y_j, x_{p_1}\} \setminus x_{p_1}) \in E(G)$. But $p < p_1$, this is a contradiction since p_1 is minimal. Consequently, $\deg_G(y_{p_1}) = 1$. Also, since G is unmixed and from Proposition 4.3 we have that $\{x_k, x_{p_1}\} = (\{x_k, x_j\} \setminus x_j) \cup (\{x_{p_1}, y_j\} \setminus y_j) \in E(G)$. Therefore, we take $w_1 = x_{p_1}$ and $w_2 = y_{p_1}$, furthermore, $e_{p_1} = \{w_1, w_2\}$. Now, we assume that $z = x_k$ and $N_G(x_k) \setminus y_k = \{y_{j_1}, ..., y_{j_t}\}$ with $k < j_1 < \cdots < j_t$. We suppose that $\deg_G(x_{j_t}) \geq 2$. If there is a vertex y_r such that $\{x_{j_t}, y_r\} \in E(G)$ then $\{x_k, y_r\} = (\{x_k, y_{j_t}\} \setminus y_{j_t}) \cup (\{y_r, x_{j_t}\} \setminus x_{j_t}) \in E(G)$, since G is unmixed and Proposition 4.3. This is a contradiction, since $r > j_t$ and j_t is maximal. Thus, there exist a vertex x_p such that $\{x_{j_t}, x_p\} \in E(G)$. But, since G is unmixed then, $\{x_k, x_p\} = (\{x_k, y_{j_t}\} \setminus y_{j_t}) \cup (\{x_p, x_{j_t}\} \setminus x_{j_t}) \in E(G)$ by Proposition 4.3. This is a contradiction, since $N_G(x_k) \setminus y_k = \{y_{j_1}, ..., y_{i_t}\}$. Hence, $\deg_G(x_{j_t}) = 1$. Therefore, we take $w_1 = y_{j_t}$ and $w_2 = x_{j_t}$, implying, $e_{j_t} = \{w_1, w_2\}$.

Finally, we assume that $z = y_k$, since y_k is not a free vertex then $N_G(y_k) \setminus x_k = \{x_{j_1}, ..., x_{j_r}\}$ with $j_1 < \cdots < j_r < k$. If y_{j_1} is not a free vertex then there is a vertex x_q such that $\{x_q, y_{j_1}\} \in E(G)$ with $q < j_1$. This implies $\{x_q, y_k\} = (\{x_q, y_{j_1}\} \setminus y_{j_1}) \cup (\{x_{j_1}, y_k\} \setminus x_{j_1}) \in E(G)$. But $q < j_1$, this is a contradiction. Therefore, $\deg_G(y_j) = 1$ and we take $w_1 = x_{j_1}$ and $w_2 = y_{j_1}$, furthermore, $e_{j_1} = \{w_1, w_2\}$.

(b) Since $e_1, ..., e_g$ is a perfect matching then there exists $i \in \{1, ..., g\}$ such that $e_i = \{z, z'\}$. Furthermore, since G is connected, $g \geq 2$ and z is a free vertex, then $\deg_G(z') \geq 2$. Hence, by incise (a) there exist $w'_1, w'_2 \in V(G)$ such that $\{z', w'_1\}, \{w'_1, w'_2\} \in E(G)$ where $\deg_G(w'_2) = 1$ and $\{w'_1, w'_2\} = e_j$ for some $j \in \{1, ..., g\}$. Therefore, we take $w_1 = z'$, $w_2 = w'_1$, $w_3 = w'_2$. Consequently $e_i = \{z, w_1\}$. $\square$

Theorem 5.4 *Let G be a graph without 3-cycles and 5-cycles. If $G_1, ..., G_k$ are the connected components of G, then the following condition are equivalent:*

(a) *G is unmixed vertex decomposable.*

(b) *G is pure shellable.*

(c) *G is Cohen-Macaulay*

(d) *G is unmixed and if G_i is not a isolated vertex then G_i has a perfect matching $e_1, ..., e_g$ of König type without squares with two e_i's.*

Proof. (a) $\Rightarrow$ (b) $\Rightarrow$ (c) It is known

(d) $\Rightarrow$ (a) Each component G_i is König then G is König. Therefore, from Proposition 5.1, G is unmixed vertex decomposable.

(c) $\Rightarrow$ (d) Since G is Cohen-Macaulay then, G is unmixed. Now, by induction on $|V(G)|$. We take $x \in V(G)$ such that $\deg_G(x)$ is minimal. If $r = \deg_G(x)$ then, we will prove that $r \leq 1$. We suppose $N_G(x) = \{z_1, ..., z_r\}$ with $r \geq 2$. Furthermore, since G does not contain 3-cycles $N_G(x)$ is a stable set. By Remark 3.16, $G' = G \setminus N_G[x]$ is a Cohen-Macaulay graph. We take $G_1', ..., G_s'$, the connected components of G'. By induction hypothesis G' satisfies (d). We can assume that $V(G_i') = \{y_i\}$ for $i \in \{1, ..., s\}$. Since $\deg_G(x)$ is minimal then, $\{y_i, z_j\} \in E(G)$ for all $i \in \{1, ..., s\}$ and $j \in \{1, ..., r\}$. If $k = k'$ then, the only maximal stable sets of G are $\{y_1, ..., y_k, x\}$ and $\{z_1, ..., z_r\}$. Thus, G is bipartite, hence G has a free vertex. This is a contradiction. Consequently, there is a component G_i' with an edge $e = \{w, w'\}$. Since $\deg_G(x)$ is minimal then, there exist $a, b \in V(G)$ such that $\{a, w\}, \{b, w\} \in E(G)$. If $a = b$ then, (a, w, w') is a 3-cycle in G. Hence, $a \neq b$. Now, if $a, b \in N_G(x)$ then (x, a, w, w', b) is a 5-cycle in G. Thus, $|\{w, w', a, b\} \cap V(G_i')| \geq 3$. Furthermore, since G_i' has a perfect matching then, $\tau(G_i') \geq 2$. Furthermore, by Corollary 5.2, G_i' has a free vertex a'. Then, by Lemma 5.3 (b), there exist edges $\{a', w_1\}, \{w_1, w_2\}, \{w_2, b'\} \in E(G_i')$ such that $\deg_{G_i'}(a') = \deg_{G_i'}(b') = 1$. By the minimality of $\deg_G(x)$ we have that a and b are adjacent with at least $r - 1$ neighbour vertices of x. If $r \geq 3$ then there exists z_j such that $z_j \in N_G(a) \cap N_G(b)$. This implies, (a, w_1, w_2, b, z_j) is a 5-cycle of G, this is a contradiction. Consequently, $r = 2$. Furthermore, we can assume that $\{a, z_1\}, \{b, z_2\} \in E(G)$, implying $C = (x, z_1, a, w_1, w_2, b, z_2)$ is a 7-cycle with $\deg_G(b) = \deg_G(a) = \deg_G(x) = 2$. Hence, by Lemma 4.9, G has C as a c-minor. This is a contradiction, since a 7-cycle is not Cohen-Macaulay and the c-minors of Cohen-Macaulay graphs are Cohen-Macaulay graphs. Therefore, $\deg_G(x) = r \leq 1$.

If $r = 0$ then, the result is clear. Now, if $r = 1$ then, $N_G(x) = \{z\}$. We can assume that $z \in V(G_1)$. Thus, the connected components of $G \setminus N_G[x]$ are $F_1, G_2, ..., G_k$ where $F_1 = G_1 \setminus N_G[x]$. By induction hypothesis $G_2, ..., G_k$ satisfy (d) and there exist $e_1 = \{x_1, y_1\}, ..., e_g = \{x_{g'}, y_{g'}\}$ a perfect matching of König type of F_1. We can assume that $D' = \{x_1, ..., x_{g'}\}$ is a minimal vertex cover of G'. Consequently, $e_1, ..., e_{g'}, e_{g'+1} = \{x, z\}$ is a perfect matching of G_1 . Furthermore, $D = \{x_1, ..., x_{g'}, z\}$ is a vertex cover of G, then $g' + 1 \leq \nu(G_1) \leq \tau(G_1) \leq |D|$. Hence, $\nu(G_1) = \tau(G_1) = g' + 1$ and $e_1, ..., e_{g'+1}$ is a perfect matching of König type of G_1. Furthermore, there are not square with two e_i's in $e_1, ..., e_{g'}$ and x is a free vertex. Therefore, G_1 does not contain square with two e_i's for $i = 1, ..., g'+1$. □

Corollary 5.5 *Let G be a graph without 3-cycles and 5-cycles. If G is Cohen-Macaulay then G has at least an extendable vertex x adjacent to a free vertex.*

Proof. From Theorem 5.4, G is König. Thus, by Corollary 5.2 there exists a free vertex x. If $N_G(x) = \{y\}$, therefore, by Remark 2.7, y is an extendable vertex. □

Definition 5.6 *Let G be a graph and let $S \subseteq V(G)$. For use below consider the graph $G \cup W_G(S)$ obtained from G by adding new vertices $\{y_i \mid x_i \in S\}$ and new edges $\{\{x_i, y_i\} \mid x_i \in S\}$. The edges $\{x_i, y_i\}$ are called whiskers (pendants). G is called whisker graph if there exists an induced subgraph H of G such that $V(H) = \{x_1, ..., x_s\}$, $V(G) = V(H) \cup \{y_1, ..., y_s\}$ and $E(G) = E(H) \cup W(H)$ where $W(H) = \{\{x_1, y_1\}, ..., \{x_s, y_s\}\}$. Its pendant edges form a perfect matching.*

Corollary 5.7 *If G is a connected graph of girth 6 or more, then the following condition are equivalent:*

(i) *G is unmixed vertex decomposable.*

(ii) *Δ_G is pure shellable.*

(iii) *$R/I(G)$ is Cohen-Macaulay.*

(iv) *G is unmixed and König.*

(v) *G is an isolated vertex or G very well-covered.*

(vi) *G is unmixed with $G \neq C_7$.*

(vii) *G is an isolated vertex or G unmixed whiskers.*

Proof. (i) ⇒ (ii) ⇒ (iii) It is known. (iii) ⇒ (iv) G is unmixed and from Theorem 5.4, G is König. (iv) ⇒ (v) From Lemma 4.4. (v) ⇒ (vi) It is clear, since C_7 is not very well-covered.

(vi) ⇒ (vii) By ([6], Corollary 5), if G is not an isolated vertex then, the pendant edges $\{x_1, y_1\}, ..., \{x_g, y_g\}$ of G form a perfect matching since $\{x_i, y_i\}$ is a pendant edge. We can assume that $\deg_G(y_i) = 1$ for all $i = 1, ..., g$. Hence, if $G' = G[x_1, ..., x_n]$ and $G = G' \cup W(V_{G'})$ with $W(V(G')) = \{y_1, ..., y_g\}$, therefore G is the whisker graph.

(vii) ⇒ (i) If G is an isolated vertex, it is clear. Now, if G is a whisker graph, then there exists a perfect matching $e_1 = \{x_1, y_1\}, ..., e_g = \{x_g, y_g\}$ such that $\deg_G(y_i) = 1$ for $i = 1, ..., g$. Thus, $D = \{x_1, ..., x_g\}$ a minimal vertex cover and $\tau(G) = g$. Hence, $e_1, ..., e_g$ is a perfect matching of König type without square with two e_i's. Therefore, by Theorem 5.4, G is unmixed vertex decomposable. □

6. Unicyclic graphs.

Definition 6.1 *A unicyclic graph is a connected graph with exactly one cycle.*

Theorem 6.2 *Let G be a unicyclic graph whose cycle is C then, the following conditions are equivalent:*

(1) *G is vertex decomposable.*

(2) *G is shellable*

(3) *G is sequentially Cohen-Macaulay.*

(4) *$|V(C)| \in \{3, 5\}$ or there exists $x \in D_1(C)$ such that $\deg_G(x) = 1$*

Proof. (1) ⇒ (2) ⇒ (3) It is known.

(3) ⇒ (4) If $|V(C)| \notin \{3, 5\}$ then, the cycle C is not sequentially Cohen-Macaulay. Furthermore, C is the only 2-connected block in G then, by Theorem 3.15 there exists $x \in D_1(C)$ such that $\deg_G(x) = 1$.

(4) ⇒ (1) If $|V(C)| = \{3, 5\}$ then, G is vertex decomposable by ([22], Theorem 1)

Now, if $|V(C)| \neq 3, 5$, then C is not vertex decomposable. Hence, by Corollary 3.17 there exists $x \in D_1(C)$ such that $\deg_G(x) = 1$. If $N_G(x) = \{y\}$ then y is a shedding vertex. Since $y \in V(C)$, then $G \setminus y$ and $G \setminus N_G[y]$ are forests, consequently they are vertex decomposable. Therefore, G is vertex decomposable. $\square$

Definition 6.3 *A clique S of a graph G containing at least one simplicial vertex of G is called a simplex of G.*

Definition 6.4 *A 4-cycle is called basic if it contains two adjacent vertices of degree two, and the remaining two vertices belong to a simplex or a basic 5-cycle of G.*

Remark 6.5 *In the case unicyclic graph we have that a 4-cycle is basic if it contains two adjacent vertices of degree two and the remaining two vertices is jointed to free vertex.*

Definition 6.6 *A graph G is in the family $\mathcal{SQC}$, if $V(G)$ can be partitioned into three disjoint subsets S_G, Q_G and C_G: the subset S_G contains all vertices of the simplexes of G, and the simplexes of G are vertex disjoint; the subset C_G consists of the vertices of the basic 5-cycles and the basic 5-cycles form a partition of C_G; the remaining set Q_G contains all vertices of degree two of the basic 4-cycles.*

Remark 6.7 *If $G \in \mathcal{SQC}$ then G is vertex decomposable by* ([11], Theorem 2.3) *and G is unmixed by* ([16], Theorem 3.1).

Lemma 6.8 *If* x *is a free vertex of* G *such that* $G' = G \setminus N_G[x]$ *is an unmixed whiskers. If there are not cycle* C *of* G *such that* $z \in V(C)$ *where* $\{x, z\} \in E(G)$ *then* G *is a whisker.*

Proof. Let $G_1, ..., G_s$ be the connected components of G then there exist induced subgraphs $H_1, ..., H_s$ such that $V(H_i) = \{x_1^i, ..., x_{r_i}^i\}$, $V(G_i) = V(H_i) \cup \{y_1^i, ..., y_{r_i}^i\}$ and $E(G_i) = E(H_i) \cup W(H_i)$ where $W(H_i) = \{\{x_1^i, y_1^i\}, ..., \{x_{r_i}^i, y_{r_i}^i\}\}$. If $\{x, z\}$ is a connected component of G then $G = \{x, z\} \cup G_1 \cup \cdots \cup G_s$ and each component is a whisker graph, therefore, G is a whisker graph. Now, if there are $w_1^i, w_2^i \in E(G_i)$ such that $\{w_1^i, z\}, \{w_2^i, z\} \in E(G)$ then there exists a path $\{w_1^i, v_1, ..., v_m, w_2^i\}$ since G_i is connected and $v_k \in V(G_i)$ for $k \in \{1, ..., m\}$, therefore, $z \in V(C) = (z, w_1^i, v_1, ..., v_m, w_2^i)$, this is a contradiction by hypothesis. Thus $|N_G(z) \cap V(G_i)| \leq 1$ for $i \in \{1, ..., s\}$. If $|V(G_i)| = 2$ then $\deg_{G_i}(x_1^i) = \deg_{G_i}(y_1^i) = 1$, therefore, if $\{w, z\} \in E(G)$ with $w \in \{x_1^i, y_1^i\}$ then $G_i \cup \{x, z\}$ is a whisker graph. We can suppose $|V(G_i)| \geq 3$, then if $\{z, y_j\} \in E(G)$ for some $j \in \{1, ..., r_i\}$ then there is a vertex w such that $\{w, x_r\} \in V(G_i)$ since $|V(G_i)| \geq 3$ then $N_{G'}(z)$ has two leaves y_j and x in $G' = G \setminus N_G[w]$, this is contradiction, therefore, $\{z, y_j\} \notin E(G)$ for all $j \in \{1, ..., r_i\}$. Thus, $\{x_j^i, z\} \in E(G)$ for some $j \in \{1, ..., r_i\}$, we can take $V(H) = \bigcup_{i=1}^s V(H_i) \cup \{z\}$, $V(G) = \bigcup_{i=1}^s V(G_i) \cup \{z, x\}$ and $E(G) = E(H) \cup W(H)$ where $W(H) = \bigcup_{i=1}^s W(H_i) \cup \{x, z\}$. Therefore, G is a whisker graph. ☐

Remark 6.9 *In* ([20], Theorem 7.3.17) *is proved that if* G *is a tree then* G *is a Cohen-Macaulay graph if and only if* G *is unmixed if and only if* G *is a whisker graph. in the following result we characterize the unicyclic graph unmixed and Cohen-Macaulay.*

Theorem 6.10 *Let* G *be a connected unicyclic graph whose cycle is* C *then,* G *is well-covered if and only if* G *satisfies one of the following conditions:*

(a) $G \in \{C_3, C_4, C_5 C_7\}$.

(b) G *is a whisker graph.*

(c) C *is a simplex 3-cycle or a basic 5-cycle and* $G \setminus V(C)$ *is a whisker graph.*

(d) C *is a basic 4-cycle with two adjacent vertices* a, b *of degree 2 in* G *such that* $G \setminus \{a, b\}$ *is a whisker graph.*

Proof. $\Rightarrow$) By induction on $l = |V(G) \setminus V(C)|$. If $l = 0$ then, $G = C$ and G satisfies (a) since the unmixed cycles are C_3, C_4, C_5 and C_7. Now, if $l \geq 1$ then, we can suppose that $C = (y_1, ..., y_k)$ with $k \geq 3$. Hence, $C \subsetneq G$. Since C is the only cycle in G then, there is a free vertex x such that $N_G(x) = \{z\}$. We can take x such that $\mathrm{d}(x, C) = max\{d(a, C) |\ a \in V(G),\ \deg_G(a) = 1\}$. Furthermore, we can suppose that $N_G(x) = \{z\}$. Since G is unmixed then, $G' = G \setminus N_G[x]$ is unmixed by Remark 3.16.

If $\mathrm{d}(x, C) \geq 2$ then $C \subseteq G'$ and G' satisfies (a), (b), (c) or (d) by induction hypothesis. If G' satisfies (a) then, $G' = C_r$ with $r \in \{3, 4, 5, 7\}$. We can assume that $C_r = (y_1, ..., y_r)$ and $\{z, y_1\} \in E(G)$. If $r = 4$ or 7 then $N_{G_2}(z)$ has two leaves x and y_1 in $G_2 = G \setminus N_G[y_3, y_{r-1}]$. This is a contradiction by Lemma 4.8, if $r = 3$ or 5. Therefore, $G'_1 = G' \setminus N_{G'}[y_2] = V(C)$ is the edge $\{x, z\}$. Then $G'_1 \cup \{x, z\}$ is a whisker graph and C_3 or C_5 is a simplex 3-cycle or a basic 5-cycle in G. Hence, G satisfies (c). If G' satisfies (b) then G is a whisker graph by Lemma 6.8. Hence, G satisfies (b). If G' satisfies (c) then C is a simplex 3-cycle or a basic 5-cycle and G'_1 is a whisker forest graph. C_3 is simplex in G since if $\{z, y_1\} \in E(G)$ and there are not simplicial vertices in C_3 then there are $z_2, z_3 \in V(G)$ such that $\{y_2, z_2\}, \{y_3, z_3\} \in E(G)$. Therefore, $N_{G_3}(z)$ has two leaves in $G_3 = G \setminus N_G[z_2, z_3]$. This is a contradiction. If C_5 is not basic in G then there are two adjacent vertices in C_5 such that $\deg_G(y_1) \geq 3$ and $\deg_G(y_2) \geq 3$ then there is z_2 such that $\{y_2, z_2\} \in E(G)$. Therefore, $N_{G_4}(z)$ has two leaves y_1 and x in $G_4 = G \setminus N_G[z_2, y_4]$. This is a contradiction. Then, C_5 is a basic 5-cycle in G. Furthermore, $G_1 = G'_1 \cup \{x, z\}$ is a whisker graph by Lemma 6.8. Thus, G satisfies (c). If G' satisfies (d). Hence, C is a basic 4-cycle (a, b, c, d) with $\deg_{G'}(a) = \deg_{G'}(b) = 2$ such that $G' \setminus \{a, b\}$ is a whisker graph. If $\{a, z\} \in E(G)$ then $N_{G_5}(z)$ has two leaves x and a in $G_5 = G \setminus N_G[c]$, this is a contradiction by Lemma 4.8. Then $\{a, z\} \notin E(G)$. Similarity, $\{b, z\} \notin E(G)$. Therefore, $\deg_G(a) = \deg_G(b) = 2$ and $G \setminus \{a, b\} = (G' \setminus \{a, b\}) \cup \{x, z\}$ is a whisker graph by Lemma 6.8. Thus, G satisfies (d).

If $\mathrm{d}(x, C) = 1$ then $V(G) = V(C) \cup D_1(C)$ and G' is a forest, therefore, G' is a whisker graph by Remark 6.9, also, there is a vertex $y_i \in V(C)$ such that $z = y_i$. We can assume $i = 1$. Since G is unicyclic, if $w \in D_1(C)$ then w is a free vertex. If each $y_i \in V(C)$ is adjacent to one free vertex then G is a whisker graph. Then G satisfies (b). If there is a vertex $y_j \in V(C)$ such that $\deg_G(y_j) = 2$, without loss of generality, we can assume $i = 2$ then, $|V(C)| \leq 5$ by ([6], Corollary 5). If $r = 3$ then C is a simplex 3-cycle. Furthermore, $G_1 = G \setminus V(C) = G \setminus N_G[y_2]$ is not a whisker graph since the vertex $\{x\}$ is isolated in G_1. Therefore, $\deg_G(y_i) = 3$ for all $i \in \{1, 2, 3\}$. Thus, G satisfies (b). Now, if $r = 5$, then $N_{G_7}(y_1)$ has two leaves x and y_2 in $G_7 = G \setminus N_G[y_4]$. This is not possible. Therefore, each $y_i \in V(C)$ is adjacent to a free vertex. Hence, G is a whisker graph. Thus, G satisfies (b). If $r = 4$, if $\deg_G(y_4) = 2$ then $N_{G'}(y_3)$ has two leaves y_2 and y_4 in G'. Consequently, $\deg_G(y_4) \geq 3$ and there is a free vertex z_4 such that $\{y_4, z_4\} \in E(G)$. Now, if $\deg_G(y_3) \geq 3$, then there is a free vertex z_3 such that $\{y_3, z_3\} \in E(G)$. Thus, $N_{G_8}(y_3)$ has two leaves y_2 and z_3 in $G_8 = G \setminus N_G[x, z_4]$. Hence, $\deg_G(y_3) = 2$ and C is a basic 4-cycle. Furthermore, $G \setminus \{y_2, y_3\}$ is a whisker graph. Therefore, G satisfies (d).

$\Leftarrow$) The families (a),(b),(c) and (d) are in $\mathcal{SQC} \cup \{C_4, C_7\}$, then by Remark 6.7 they are unmixed. $\square$

Corollary 6.11 *Let G be an unicyclic graph then, the following condition are equivalent:*

(1) *G is unmixed vertex decomposable.*

(2) *G is pure shellable*

(3) *G is Cohen-Macaulay.*

(4) *G is unmixed and $G \neq C_4, C_7$.*

Proof. (1)⇔(2)⇔(3) By Theorem 6.2.

(3) ⇒ (4) Since G is Cohen-Macaulay then G is unmixed. Furthermore, C_4 and C_7 are not Cohen-Macaulay.

(4) ⇒ (1) Let C be the cycle of G. By Theorem 6.10, $G \in \mathcal{SQC}$ since $G \neq C_4, C_7$. Therefore G is vertex decomposable by Remark 6.7. $\square$

Corollary 6.12 *If G is an unmixed unicyclic then G is vertex decomposable if and only if $G \neq C_4, C_7$.*

Proof. By Corollary 6.11. $\square$

Referencias

[1] J. Biermann, C. A. Francisco, H. Tài Hà, A. Van Tuyl, *Colorings of simplicial complexes and vertex decomposability*. Preprint, 2012, arXiv:1209.3008v1.

[2] A. Bruns, J. Herzog, *Cohen-Macaulay Ring*. Cambridge University Press, Cambridge, 1997.

[3] I. Castrillón, R. Cruz, *Escalonabilidad de grafos e hipergrafos simples que contienen vértices simpliciales*. Matemáticas: Enseñanza Universitaria, Vol. XX, (2012), No. 1, 69-80.

[4] M. Crupi; G. Rinaldo; N. Terai, *Cohen-Macaulay edges ideals whose height is half of the number of vertices*. Nagoya Math. J., **201** (2011), 116-130.

[5] M. Estrada and R. H. Villarreal, *Cohen-Macaulay bipartite graphs*. Arch. Math. **68** (1997), 124-128.

[6] A. Finbow, B. Hartnell, R.J. Nowakowski, *A characterization of well covered graphs of girth 5 or greater*. Journal of Combinatorial Theory. **B 57** (1993), 44-68.

[7] A. Finbow, B. Hartnell, R.J. Nowakowski, *A characterization of well covered graphs that contain neither 4-nor 5-cycles*. Journal of Graph Theory. **18** (1994), 713-721.

[8] C. A. Francisco, A. Van Tuyl, *Sequentially Cohen-Macaulay edges ideals*. Proc. Amer. Math. Soc. (electronic) **135** (2007), no 8, 2327-2337.

[9] I. Gitler, C. E. Valencia, *On bounds for some graph invariants*. Boletín de la Sociedad Matemática Mexicana 10 (2005).

[10] J. Herzog; T. Hibi, *Distributive lattices, bipartite graphs and Alexander duality*. J. Alg. Combin. **22** (2005), 289-302.

[11] D. T. Hoang; N. C. Minh; T.N. Trung, *On then Cohen-Macaulay graph and girth*. Preprint 2012 arxiv: 1204.5561v2.

[12] M. Katzman, *Characteric-independence of Betti numbers of graph ideals*. J. Combin. Theory Ser. **A 113** (2006) 435-454.

[13] M. Mahmoudi; A. Mousivand; M. Crupi; G. Rinaldo; N. Terai; S. Yassemi, *Vertex decomposability and regularity of very well-covered graphs*. Journal of Pure and Applied Algebra **215** (2011), 2473-2480.

[14] S. Morey, E. Reyes, R.H. Villarreal, *Cohen-Macaulay, shellable and unmixed clutters with perfect matching of König type*. Journal of Pure and Applied Algebra **212** (2008), 1770-1786.

[15] E. Prisner, J. Topp, P.D. Vestergaard, *Well covered simplicial, chordal and circular arc graphs*. Journal of Graph Theory. **21-2** (1996), 113-119.

[16] B. Randerath, L. Volkmann, *A characterization of well covered block-cactus graphs*. Australas. Journal of Combinatorics. **9** (1994), 307-314.

[17] R. P. Stanley, *Combinatorics and Commutative Algebra. Second edition*. Progress in Mathematics 41. Birkhäuser Boston, Inc., Boston, MA, 1996.

[18] J. Topp and L. Volkmann, Characterization of unicyclic graphs with equal domination and independence number. Discuss. Math, **11** (1991), 27-34.

[19] A. Van Tuyl; R. H. Villarreal, *Shellable graphs and sequentially Cohen-Macaulay bipartite graphs*, Journal of Combinatorial Theory Series **A 115** (2008), 799-814. ISSN:0097-3165

[20] R. H. Villarreal, *Monomial Algebras*. Second edition, Monographs and Research Notes in Mathematics, Chapman and Hall/CRC, 2015.

[21] M. L. Wachs, *Obstructions to shellability* Discrete Comput. Geom. 22 (1999), no 1, 95-103.

[22] R. Woodroofe, *Vertex decomposable graphs and obstruction to shellability*. Proc. Amer. Math. Soc. **137**, (2009), 3235-3246.

Departamento de Matemáticas, Centro de Investigación y de Estudios Avanzados del IPN, Apartado Postal 14-740, 07000 Mexico City, D.F.
E-mail address: `idcastrillon@math.cinvestav.mx`

Instituto de Matematicas, Universidad de Antioquia, Apartado Aéreo 1226, Medellín, Colombia.
E-mail address: `roberto.cruz@udea.edu.co`

Departamento de Matemáticas, Centro de Investigación y de Estudios Avanzados del IPN, Apartado Postal 14-740, 07000 Mexico City, D.F.
E-mail address: `ereyes@math.cinvestav.mx`